
Rate Equations Map

Cambridge International AS & A Level Chemistry (9701) · Topic 26 Reaction kinetics · Rate equations

1 Measuring a rate

26.1.2

- Rate = change in concentration ÷ time
- Unit: **mol dm^{-3} s^{-1}**
- Gas volume, mass loss, **colorimetry**
- Titrate **quenched** samples



2 Rate equations

26.1.1-26.1.2


- rate = **k[A]^m[B]^n**
- Orders 0, 1 or 2: **from experiments**
- Overall order = m + n
- k fixed at one temperature




3 Units of k

26.1.4

- Zero: mol dm^{-3} s^{-1}
- First: **s^{-1}**
- Second: dm^3 mol^{-1} s^{-1}
- Third: dm^6 mol^{-2} s^{-1}






4 Initial rates

26.1.2


- Change **one** concentration only
- Double it: rate × 1, × 2, × 4
- Clock reaction: rate ∝ **1/t**
- Same total volume: water



5 Graph shapes

26.1.2


- [A]–time: zero is a **straight line**
- Rate = gradient of the **tangent**
- Rate–[A]: flat, straight, curve
- First order: gradient = k



6 Half-life

26.1.3-26.1.4

- Time for [A] to **halve**
- **Constant** t½: first order
- **k = 0.693/t½**
- Second order: t½ doubles



Orders at a glance

one reactant A; doubling [A] at constant temperature

Order	Rate equation	Unit of k	[A]–time graph	Rate–[A] graph	Double [A]
0	rate = k	mol dm^{-3} s^{-1}	straight line	horizontal line	× 1
1	rate = k[A]	s^{-1}	curve, constant t½	straight, through 0	× 2
2	rate = k[A]^2	dm^3 mol^{-1} s^{-1}	curve, t½ doubles	upward curve	× 4

Worked examples

Order from a table

- [B] × 3, rest the same
- rate × 9 = 3^2
- **second order** in B

k from a half-life

- t½ = 139 s each time
- k = 0.693/139
- **5.0 × 10^{-3} s^{-1}**

Clock times

- [I$^-$] × 2: t 48 s → 24 s
- 1/t doubles
- **first order** in I$^-$

Rate equations lead on to the rate-determining step, temperature and catalysts (Topic 26 Mechanisms, temperature and catalysts).

Online Learning System · onlinelearningsystem.net · © 2026 Online Learning System. All rights reserved. Not for redistribution.