Redox Titrations Map

Edexcel International A Level Chemistry · Unit 5 · Topic 16B Redox Titrations


1 Manganate(VII) and iron(II) 16.15

- MnO_4^- + $8H^+$ + $5Fe^{2+}$: **1 : 5**
- Purple → colourless
- **Dilute H_2SO_4** only
- Not HCl, HNO_3 or ethanoic acid



2 Core Practical 13a 16.17


- Pipette Fe^{2+}, excess acid
- **Self-indicating**
- End: first permanent **pale pink**
- Read the **top** of the meniscus



3 Other reducing agents 16.15

- $C_2O_4^{2-}$: **2 : 5**, warm to 60 °C
- Mn^{2+} catalyses: autocatalysis
- H_2O_2 → O_2: **2 : 5**
- FeC_2O_4: **3 : 5**





4 Iodine and thiosulfate 16.15, 16.17

- Oxidant + **excess KI** → I_2
- $I_2 + 2S_2O_3^{2-} \rightarrow 2I^- + S_4O_6^{2-}$
- Starch near the end: **blue-black**
- End: **colourless**




5 Ratios through iodine 16.15

- Cu^{2+} : $S_2O_3^{2-}$ = **1 : 1**
- ClO^- : $S_2O_3^{2-}$ = **1 : 2**
- IO_3^- : $S_2O_3^{2-}$ = **1 : 6**
- Moles, ratio, scale up



6 Uncertainty and validity 16.16

- Titre ±0.10 cm^3 (two readings)
- % = uncertainty ÷ value × 100
- **Add** the percentages
- % difference > % uncertainty: **systematic**



Reacting ratios in redox titrations

titrant : substance

Titrant	Substance	Ratio
MnO_4^-	Fe^{2+}	1 : 5
MnO_4^-	$C_2O_4^{2-}$ or H_2O_2	2 : 5
MnO_4^-	FeC_2O_4	3 : 5
$S_2O_3^{2-}$	I_2	2 : 1
$S_2O_3^{2-}$	Cu^{2+} (via I_2)	1 : 1
$S_2O_3^{2-}$	IO_3^- (via I_2)	6 : 1

Worked examples

Iron tablet

- 11.20 cm^3 of 0.00500 MnO_4^-
- × 5 × 10 × 55.8
- **0.156 g Fe**

Brass

- 18.90 cm^3 of 0.100 $S_2O_3^{2-}$
- 1 : 1, × 10 × 63.5
- **1.20 g Cu**

Uncertainty

- 22.30 ± 0.10 cm^3
- 0.10 ÷ 22.30 × 100
- **0.45%**

Electrode potentials explain the choice of acid (Topic 16A); Mn^{2+} autocatalysis returns in Topic 17.

Online Learning System · onlinelearningsystem.net · © 2026 Online Learning System. All rights reserved. Not for redistribution.