

Ionic Bonding Map

Edexcel A Level Chemistry · Topic 2 Bonding and Structure · Ionic Bonding

1 Forming ions

2A.3

- Metals **lose** electrons: positive ions (cations)
- Non-metals **gain** electrons: negative ions (anions)
- Charge = electrons lost or gained
- Often a noble-gas configuration: Na^+ 2,8; Cl^- 2,8,8
- Groups 1–3: 1+, 2+, 3+; Groups 5–7: 3–, 2–, 1–

2 Dot-and-cross diagrams

2A.4

- Outer shells only; dots for one atom, crosses for the other
- Cation: its new outer shell (8) or drawn empty
- Anion: its own electrons + those gained
- Square brackets** and the **charge** on every ion
- Ratio from electron transfer: CaF_2 , Na_2O , Al_2O_3



3 The giant lattice

2A.1, 2B.23

- A **giant lattice**: regular 3D array of ions
- NaCl : each ion has **six** opposite neighbours
- Ionic bonding: strong **net electrostatic attraction**
- Acts in all directions: no molecules
- The formula is only the ratio of ions

4 Bond strength

2A.2

- Higher charges**: stronger attraction
- Smaller ions**: centres closer, stronger attraction
- Stronger bonding: higher melting temperature
- MgO 2852 °C against NaCl 801 °C
- $\text{NaF} > \text{NaCl} > \text{NaBr} > \text{NaI}$ (anion grows)



5 Ionic radii

2A.5

- Larger **down a group**: one more shell
- Cations smaller than their atoms; anions larger
- Isoelectronic**: same electrons, more protons, smaller
- N^{3-} 146 > O^{2-} > F^- > Na^+ > Mg^{2+} > Al^{3+} 54 pm
- The most negative ion of a set is largest



6 Radii and bond strength

2A.2, 2A.5

- Ion separation = $r_+ + r_-$
- Closer ions: **stronger** attraction
- Same charges: the separation decides ($\text{LiF} > \text{KF}$)
- Charges matter most: MgO 2852 °C, NaF 993 °C
- Both sets of ions have ten electrons

Charge, radius and melting temperature

radii are ionic radii in pm; melting temperatures in °C

Compound	Ion charges	Cation radius	Anion radius	Melting point
NaF	1+, 1–	102	133	993
NaCl	1+, 1–	102	181	801
KCl	1+, 1–	138	181	770
NaI	1+, 1–	102	220	661
CaO	2+, 2–	100	140	2613
MgO	2+, 2–	72	140	2852

Evidence for ions (2A.6)

Melting and brittleness

- High melting temperatures
- Hard: strong attraction in all directions
- Brittle: like charges meet and repel

Conduction and solubility

- Conduct only when molten or dissolved
- Decomposed at the electrodes
- Polar water hydrates the ions

Migration of ions

- Copper(II) chromate(VI) on moist paper
- Blue Cu^{2+} moves to the – electrode
- Yellow CrO_4^{2-} moves to the + electrode

Ionic and covalent bonding are the two ends of a continuum (Covalent Bonding); lattice energy measures ionic bonding (Topic 13).



Online Learning System · onlinelearningsystem.net · © 2026 Online Learning System. All rights reserved. Not for redistribution.